

11-20-2015

Cardiopulmonary Exercise Testing-Based Algorithm and Its Usefulness in Clinical Cardiology

Mirza Nubair Ahmad

Syed Hasan Yusuf

Rafath Ullah


Mary Ellis

Haroon Yousaf

Timothy E. Paterick

Khawaja Afzal Ammar

Follow this and additional works at: <https://aurora.org/jpcrr>

 Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), and the [Other Analytical, Diagnostic and Therapeutic Techniques and Equipment Commons](#)

Recommended Citation

Ahmad M, Yusuf S, Ullah R, Ellis M, Yousaf H, Paterick TE, Ammar KA. Cardiopulmonary exercise testing-based algorithm and its usefulness in clinical cardiology. *J Patient Cent Res Rev*. 2015;2:204-205. doi: 10.17294/2330-0698.1223

Journal of Patient-Centered Research and Reviews (JPCRR) is a peer-reviewed scientific journal whose mission is to communicate clinical and bench research findings, with the goal of improving the quality of human health, the care of the individual patient, and the care of populations.



worker consultation and diagnoses of respiratory issues and dementia.

Conclusion: A promising automated model generated by EHR data to predict 30-day readmissions and mortality among hospitalized older adults, these findings will be used by the health care system to incorporate a real-time alert into physician workflow. Efforts to improve care will include interventions targeted at the highest-risk group.

FIRST PLACE POSTER (tie)

See page 217 for citation.

SECOND PLACE POSTER

See page 217 for citation.

THIRD PLACE POSTER

Echocardiographic Predictors of Admission Among Patients With Heart Failure With Reduced Ejection Fraction

Chi C. Cho, Yang Shi, Robyn Shearer, Nasir Z. Sulemanjee, Dianne L. Zwicke, T. Edward Hastings, Omar M. Cheema, Vinay Thohan

Aurora Research Institute, Aurora Health Care; Aurora Cardiovascular Services, Aurora Health Care

Background: Congestive heart failure afflicts 5.7 million people in the United States with annual incidence of 600,000 and mortality of 280,000. Heart failure accounts for greater than 1 million hospitalizations annually and the single largest inpatient Medicare expense. As the U.S. population ages and greater emphasis is placed on population health as a means to bend projected health care expenditures, large health care organizations will need to develop algorithms to identify patients at high risk with heart failure and possibly preempt hospitalizations. Doppler echocardiography is routinely performed in clinical assessment of severe heart failure.

Purpose: We sought to determine echocardiographic parameters that predict 1-year cardiac events among ambulatory patients diagnosed with heart failure with reduced ejection fraction.

Methods: A retrospective single-institution investigation identified 485 patients aged < 75 years with left ventricular ejection fraction < 35%. Kaplan-Meier method was used to identify parameters that corresponded with primary endpoint of hospitalization, emergency room visit or death.

Results: High risk of primary endpoint could be segregated into four groups by presence of one or more of the following parameters (0, 1, 2, 3): mitral inflow E/A ratio > 1.5, mitral E-wave deceleration time < 160 ms or peak tricuspid regurgitant (TR) velocity > 3 m/s. Event-free survival was significantly lower in high-risk group compared to low-risk group (P=0.002). The 30-day hospitalization rates among those with all three factors compared to those without was 37.5% and 17.3%, P=0.018.

Conclusion: Presence of routine echocardiographic parameters, including E/A ratio > 1.5, E-wave deceleration time < 160 ms and TR velocity > 3 m/s, is associated with high cardiovascular event rates among nonhospitalized ambulatory patients with reduced ejection fraction heart failure.

RIESELBACH DISTINGUISHED PAPER #1

Cardiopulmonary Exercise Testing-Based Algorithm and Its Usefulness in Clinical Cardiology

Mirza Nubair Ahmad, Syed Hasan Yusuf, Rafath Ullah, Mary Ellis, Haroon Yousaf, Timothy E. Paterick, Khawaja Afzal Ammar

Aurora Cardiovascular Services, Aurora Health Care; Division of Pulmonary and Critical Care Medicine, Medical College of Wisconsin

Background: Only cardiopulmonary exercise (CPX) testing provides information on the ability of the cardiovascular system to meet the body's metabolic demands in terms of oxygen consumption (VO_2) and carbon dioxide production (VCO_2). However, CPX testing is underutilized by cardiologists due to complex diagnostic algorithms involving up to 30 variables as well as lack of validation studies. In addition, CPX also provides oxygen (O_2) pulse as a continuous measure of stroke volume, which is its superiority to other stress modalities in which systolic function is measured at peak stress and rest. In the literature, it has been recommended that a composite criterion (combining peak O_2 pulse with O_2 pulse curve pattern) should be used to assess the cardiac function. Furthermore, the operating test characteristics and optimal cutoff of O_2 pulse for distinguishing cardiac from noncardiac causes of exercise limitation also are unknown.

Purpose: We tested whether a 6-variable algorithm would discriminate cardiac from noncardiac causes of dyspnea when compared with comprehensive CPX testing to promote its use by cardiologists. We also tested several cutoff points along with the composite criterion against the clinical standard to define the optimal O_2 pulse cutoff point.

Methods: Consecutive patients (n=54) referred for dyspnea underwent CPX test consisting of pulmonary (VO_2 , VCO_2 , 22 additional variables and invasive measurement of lactate and blood gases at peak and baseline) and cardiac (exercise ECG, heart rate and blood pressure response) components as well as medical record evaluation. Patients were categorized as normal or abnormal by an experienced pulmonologist. Abnormal patients were further categorized according to cause of dyspnea (cardiac, pulmonary, deconditioning, poor effort and miscellaneous). Subsequently, the 6-variable algorithm was applied by a cardiologist blinded to all of the information from CPX tests, and the patients were categorized similarly. The 6 variables used were peak O_2 uptake, peak respiratory exchange ratio, O_2 pulse, heart rate reserve, breathing reserve ($1 - [\text{peak ventilation (VE) / maximal voluntary ventilation}]$) and ventilatory efficiency (VE/VCO_2). Seven O_2 pulse reference cutoff points

included nongender-based (<15 ml/beat), gender-based (<15 ml/beat for males and <10 ml/beat for females) and < 80% of O₂ pulse based on five different definitions of predicted VO₂ max. The optimal cutoff obtained was then used to create the composite criterion. For the purpose of evaluating this composite criterion, the study population was recategorized as: noncardiac group (n=18), normal patients according to the composite criterion; or cardiac group (n=13), abnormal patients according to the composite criterion. Patients who were normal by only one component of the composite criterion were categorized as borderline (n=23). Data were analyzed against the comprehensive CPX test by first excluding the borderline patients and then by including them with either the cardiac or noncardiac group.

Results: The 6-variable algorithm performed well against comprehensive CPX test in discriminating cardiac from noncardiac causes of dyspnea, with 94% sensitivity, 92% specificity, 84% positive predictive value (PPV), 97% negative predictive value (NPV) and 93% accuracy. The results remained consistent for gender and referral source. O₂ pulse, as defined by Wasserman, had the highest accuracy, specificity and PPV and therefore was used to define the composite criterion. The composite criterion had an accuracy of 87%, PPV of 77%, NPV of 94%, sensitivity of 91% and specificity of 85%, when borderline patients were excluded. Including borderline patients in the cardiac group (n=36) improved sensitivity (94%) and maintained NPV (94%) but greatly decreased specificity (46%), PPV (44%) and accuracy (61%), whereas including these patients in the noncardiac group (n=41) improved specificity (92%) and maintained similar PPV (77%) and accuracy (81%) but decreased sensitivity (59%) and NPV (83%).

Conclusion: This is the first study to validate a diagnostic algorithm for patients undergoing CPX testing as well as demonstrate that a simplified 6-variable algorithm applied by a cardiologist without prior CPX experience is quite accurate to evaluate the optimal O₂ pulse value at peak stress for discrimination of cardiac and noncardiac causes, and to provide the operating test characteristics for the common clinical practice of using composite criterion to diagnose cardiac versus noncardiac causes of dyspnea.

RIESELBACH DISTINGUISHED PAPERS #2-4

See page 217 for citations.

SELECT ABSTRACTS

Score Big for Decreasing Mortality: ICD Risk Score Model

Linda Francaviglia, Rachel Petersen, Maria Stone, M. Eymann Mortada

Departments of Cardiovascular Data Services and Quality Management, Aurora Cardiovascular Services, Aurora Health Care

Background: Aurora Health Care, a system of 14 acute care hospitals in eastern Wisconsin, has been a long-time

participant in the American College of Cardiology's National Cardiovascular Data Registries, submitting data to its ICD Registry™ since 2005. Our system's implantable cardioverter-defibrillator (ICD) procedure volume averages 930 cases annually. During 2012 we experienced an increase in in-hospital mortality/morbidity for ICD cases.

Purpose: A single-center study examining in-hospital mortality/morbidity post-ICD implant before and after changes in practice and patient selection.

Methods: ICD implants and generator changes discharged from January 1, 2009, to December 31, 2012, were included in developing a risk model predicting in-hospital mortality/morbidity. The risk score was shared with physicians for clinical input. A point system was developed, including those factors with highest risk. Using the defined factors, a risk score > 14 was used to indicate those at highest risk for morbidity/mortality. The risk score model was fit on the development group (2009–2012), and then re-run for the intervention cohort from January 1, 2013, to June 30, 2014. Logistic regression was used in the risk model development and validation. Continuous variables were compared using Student's t-test, and categorical variables were compared using chi-square test.

Results: From 2009 to 2012, 3,417 ICD implants and generator changes were performed and included in risk model development. Of those, 200 (5.9%) patients were indicated as high risk with a score > 14. From January 2013 to June 2014, 1,057 implants and generator changes were performed, with 41 (3.4%) patients indicated as high risk with a score > 14. In the development phase, mean age was 67 years and 70% of patients were male. Post-model development, mean age was 66 years with 72% male. For patients indicated as high risk, in-hospital mortality/morbidity dropped from 20 (10%) to 2 (4.9%), though the decrease was not statistically significant (P=0.39).

Conclusion: Awareness of high-risk patients and changes in patient selection can lead to improvement in in-hospital mortality/morbidity among those high-risk patients. Although the improvement was not statistically significant, this was most likely due to low volumes and we will continue to monitor outcomes among these patients.

Geographic Distribution of Infant Death During Birth Hospitalization and Maternal Group B *Streptococcus* Colonization: Eastern Wisconsin

Jessica J.F. Kram, Dennis J. Baumgardner, Kiley A. Bernhard, Melissa A. Lemke

Center for Urban Population Health; Department of Family Medicine, Aurora UW Medical Group; TRIUMPH, University of Wisconsin-Madison

Background: Neonatal death rate in the United States is 4/1,000 live births; infant death rate is 6/1,000. Group B *Streptococcus* (GBS) may be transmitted from a colonized mother (rates vary from 15% to 35%) to the newborn during a vaginal delivery, and may contribute to neonatal death.